



US007574340B2

(12) **United States Patent**  
**Bavari et al.**

(10) **Patent No.:** **US 7,574,340 B2**  
(45) **Date of Patent:** **Aug. 11, 2009**

(54) **SMALL MOLECULES AND A  
PHARMACOPHORE MODEL FOR  
INHIBITION OF BOTULINUM TOXIN AND  
METHODS OF MAKING AND USING  
THEREOF**

(75) Inventors: **Sina Bavari**, Frederick, MD (US);  
**James J. Schmidt**, Frederick, MD (US);  
**James Burnett**, Richmond, VA (US);  
**Rick Gussio**, Frederick, MD (US)

(73) Assignee: **The United States of America as  
represented by the Secretary of the  
Army**, Washington, DC (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 219 days.

(21) Appl. No.: **10/935,622**

(22) Filed: **Sep. 8, 2004**

(65) **Prior Publication Data**

US 2005/0153945 A1 Jul. 14, 2005

**Related U.S. Application Data**

(60) Provisional application No. 60/501,243, filed on Sep.  
8, 2003.

(51) **Int. Cl.**  
**G06G 7/48** (2006.01)

(52) **U.S. Cl.** ..... **703/11**

(58) **Field of Classification Search** ..... 702/19;  
703/11

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

6,531,126 B2 3/2003 Farmer ..... 424/115  
2005/0038246 A1\* 2/2005 Arnold et al. .... 544/262

**OTHER PUBLICATIONS**

Lacey et al. "Crystal Structure of Botulinum Neurotoxin Type A and  
Implications for Toxicity," Nature Structural Biology (1998) vol. 5,  
No. 10, pp. 898-902.\*

Schmidt et al. "Type A Botulinum Neurotoxin Proteolytic Activity:  
Development of Competitive Inhibitors and Implications for Sub-  
strate Specificity at the S1' Binding Subsite," FEBS Letters (1998)  
vol. 435, pp. 61-64.\*

Deshpande et al. "Efficacy of Certain Quinolines as Pharmacological  
Antagonists in Botulinum Neurotoxin Poisoning," Toxicon (1997)  
vol. 35, No. 2, pp. 433-445.\*

Penzotti et al. "A Computational Ensemble Pharmacophore Model  
for Identifying Substrates of P-Glycoprotein," Journal of Medicinal  
Chemistry (2002) vol. 45, No. 9, pp. 1737-1740.\*

Bajorath, Jurgen "Integration of Virtual and High-Throughput  
Screening," Nature Reviews Drug Discovery (2002) vol. 1, pp. 882-  
894.\*

Vennerstrom, et al. (1998) "Bisquinolines. 2. Antimalarial N,N-  
Bis(7-chloroquinolin-4-yl)heteroalkanediamines" J. Med. Chem.  
41:4360-4364.

\* cited by examiner

*Primary Examiner*—Jerry Lin

(74) *Attorney, Agent, or Firm*—Elizabeth Arwine

(57) **ABSTRACT**

Disclosed herein is a pharmacophore model for inhibiting  
Botulinum neurotoxin A metalloprotease activity which com-  
prises a first plane A, a second plane B, a first hydrophobic  
moiety C, a second hydrophobic moiety D and a positive  
ionizable substituent E. The pharmacophore model may fur-  
ther comprise a heteroatom in the first plane A. In some  
embodiments, the distance between the center of the first  
plane A and the center of the second plane B is about 6.5 to  
about 9.5 Å. In some embodiments, the distance between the  
center of the first hydrophobic moiety C and the center of the  
second hydrophobic moiety D is about 8.0 to about 16.0 Å. In  
some embodiments, the distance between the center of the  
first plane to the center of the first hydrophobic moiety C is  
about 3.0 to about 5.0 Å. In some embodiments, the distance  
between the center of the second plane to the center of the  
second hydrophobic moiety C is about 3.0 to about 5.0 Å. In  
some embodiments, the distance between the center of the  
first plane to the center of the positive ionizable substituent is  
about 6.5 to about 9.5 Å.

**17 Claims, 10 Drawing Sheets**